



BEQVEZ[™]

fidanacogene elaparvovec-dzkt

Suspension for intravenous infusion 1×10^{13} vg/mL

What to expect on the BEQVEZ treatment journey

Bill, who received the BEQVEZ infusion in 2016*

*Bill participated in the BEQVEZ clinical study. He was compensated for sharing his story.



This guide gives you:

- Information about the steps in your BEQVEZ treatment journey
- Guidance on how to get educational resources
- Contact information for support services available to you

Always consult your doctor for treatment-related questions. This brochure is not intended to replace the advice of your care team.

What is BEQVEZ?

BEQVEZ is a one-time gene therapy used for the treatment of adults with moderate to severe hemophilia B who are receiving routine prophylaxis, or have a current life-threatening bleed or a history of life-threatening bleeds, or have repeated serious spontaneous bleeds.

Before treatment with BEQVEZ, your healthcare professional will conduct a blood test to check for neutralizing antibodies to the AAVRh74var virus. The results of this testing will help determine if you may receive BEQVEZ.

Important Safety Information

Before receiving BEQVEZ, tell your healthcare professional about all your medical conditions, including if you:

- Have liver problems, including hepatitis
- Have factor IX inhibitors or a history of factor IX inhibitors
- Have active HIV infection

Please see Important Safety Information throughout and full [Prescribing Information](https://www.BEQVEZ.com) at www.BEQVEZ.com.

What to expect if you are considering gene therapy with BEQVEZ

You may have made the important decision to see if BEQVEZ is the right treatment choice for you. We're here to support you at every step of your BEQVEZ treatment journey.

It is important to remember that individual results and duration of effect may vary. It is not possible to predict who will respond and how long the treatment response will continue. After receiving the BEQVEZ infusion, regular post-infusion monitoring is required and it is important to keep factor IX (FIX) on hand.



Confirm your eligibility

To find out if BEQVEZ is right for you, you and your doctor will talk about considerations like potential lifestyle changes. You will also need lab testing and confirmation of your access and insurance coverage to see if you're eligible for BEQVEZ.

Once your overall eligibility is confirmed, your care team will give you information about the next step on your BEQVEZ journey.

The infusion center will provide you with information on what you can expect on the day of the infusion. This includes the schedule for the day, lab work needed, and estimated time of discharge.

How do I find out if BEQVEZ is the right treatment option for me?

Your doctor will check your eligibility with various tests to confirm if BEQVEZ may be a treatment option for you. These tests include, but are not limited to, checking your liver health, checking for FIX inhibitors, and seeing if you have any neutralizing antibodies (NABs) to BEQVEZ. Your care team will talk to you about your results and whether BEQVEZ is right for you.

You will get the results from your NAb test after about 7 to 10 days.

What happens if I have NABs to BEQVEZ?

Your doctor can help by discussing your hemophilia B treatment options if BEQVEZ is not right for you.

How can I get help understanding my insurance coverage for BEQVEZ?

Pfizer GeneTogether is here to help provide personalized support throughout your BEQVEZ journey.

The program includes a dedicated **Patient Case Manager**. They can help you understand your health insurance coverage and out-of-pocket costs. They can also help eligible commercially insured patients understand their co-pay support options.

Pfizer
genetogether™

To get started and see the support available:

Call us at 1-855-4BEQVEZ (1-855-423-7839)

9:00 AM to 7:00 PM ET Monday to Friday

Visit us at [pfizergenetogether.com](https://www.pfizergenetogether.com)

Please see Important Safety Information throughout and full [Prescribing Information](https://www.BEQVEZ.com) at www.BEQVEZ.com.



Preparing for the infusion

How do I know where to go for my BEQVEZ infusion?

The infusion happens at a gene therapy infusion center. Your care team can direct you to the right infusion center.

Can I get help arranging travel to the center?

For commercially insured eligible patients, your Patient Case Manager can assist with travel logistics required for your treatment.



On the infusion day

What can I expect on the day of the infusion?

On the day of your infusion, you should plan to spend all day at the gene therapy infusion center. The infusion takes about 1 hour, but you will need to be at the center before and after the infusion.

The process before, during, and after your infusion may vary across centers. The infusion center you use will provide you with information to guide you through each part of the infusion process.



During the infusion

How long will the infusion take and can my loved ones be with me?

The infusion takes about 1 hour. You may want to bring something to read or music to listen to. You may also be able to have loved ones with you during the infusion. Ask your care team if this is possible at the center you go to.

What are the possible side effects or risks of BEQVEZ?

In the clinical trials of BEQVEZ, the most common side effect was increased liver enzymes. These cases were treated with corticosteroids. These are not all the possible side effects of BEQVEZ. Your care team will monitor you for possible infusion reactions on the day of your infusion.

Please read the Important Safety Information throughout this guide.

THE INFUSION TOOK ABOUT AN HOUR.

- Bill

Always consult your doctor for treatment-related questions.

Important Safety Information (cont'd)

BEQVEZ may cause serious side effects, including:

Increased Liver Enzymes. Most patients treated with BEQVEZ developed elevated liver enzyme levels and most did not experience any symptoms.

Your healthcare professional will **monitor liver enzymes and factor IX activity levels** before administration of BEQVEZ and frequently following the administration to detect and identify possible elevations in liver enzymes and to monitor your response to BEQVEZ. Your doctor may prescribe a corticosteroid for the treatment of elevated liver enzymes.

Please see Important Safety Information throughout and full **Prescribing Information** at www.BEQVEZ.com.


BEQVEZ[™]
fidanacogene elaparvovec-dzkt
Suspension for intravenous infusion 1×10^{13} vg/mL



What to expect after the infusion and in your scheduled follow-ups



After the infusion

How will I know what to do after the infusion?

After the infusion, you'll stay at the infusion center for at least 3 hours for monitoring. Your care team will:

- Tell you when you are ready to leave the infusion center
- Provide you with a post-infusion monitoring plan

It will be important to see your care team for post-infusion monitoring of your liver and FIX activity levels routinely.



After-treatment experience

What does my care team check in the post-infusion follow-ups?

You will have blood tests to assess your liver health and FIX activity levels.

In the clinical trial for BEQVEZ, some participants had raised liver enzyme levels and/or a decrease in FIX activity levels. Depending on their levels, these participants were prescribed corticosteroids.

What lifestyle considerations should I know about before receiving BEQVEZ?

You may need to make temporary changes to your lifestyle. Your doctor will talk to you about:

- Limiting alcohol during the first year after the BEQVEZ infusion
- Not donating sperm and being abstinent or using a male condom for the first 6 months after your infusion

Please see Important Safety Information throughout and full [Prescribing Information](https://www.BEQVEZ.com) at www.BEQVEZ.com.

After-treatment experience (continued)

After the infusion, when do I need to have scheduled follow-ups?

Your care team will help you plan for scheduled follow-ups.

Post-infusion follow-ups, which include blood draws, are done 1 or 2 times a week in the first 4 months after receiving BEQVEZ and lessen over time.



THE CHECK-INS
LESSEned
OVER TIME.

- Bill



Check out Bill's story
at [BEQVEZ.com](https://www.BEQVEZ.com) and sign
up for educational emails

Important Safety Information (cont'd)

Limit alcohol consumption during the first year following BEQVEZ infusion, as alcohol may reduce the effect of BEQVEZ and may increase liver enzyme levels.

Infusion reactions, including hypersensitivity and severe allergic reactions (anaphylaxis) may occur. Alert your healthcare professional right away if you get any symptoms of hypersensitivity, which may include but are not limited to low blood pressure, fever, heart palpitation, nausea, vomiting, chills, or headache.

BEQVEZ can insert itself into the DNA of cells in the human body. The effect that insertion may have on those cells is unknown but may contribute to a theoretical risk of cancer. There have been no reported cases of cancer caused by treatment with BEQVEZ.

The most common side effect of BEQVEZ is increased liver enzymes. These are not all the possible side effects of BEQVEZ. For more information, ask your healthcare professional.

Please see Important Safety Information throughout
and full [Prescribing Information](https://www.BEQVEZ.com) at www.BEQVEZ.com.


BEQVEZTM
fidanacogene elaparvovec-dzkt
Suspension for intravenous infusion 1×10¹³vg/mL



Call 1-855-4BEQVEZ (1-855-423-7839)
9:00 AM to 7:00 PM ET Monday to Friday

- Get one-on-one patient support and information
- Connect with a Pfizer Patient Affairs Liaison (PAL)* to help you access the educational support and information you need



Visit [BEQVEZ.com](https://www.BEQVEZ.com)
to learn more

Pfizer has over **25 years** of commitment
to the hemophilia community

*Pfizer PALs are not measured or awarded based on sales performance nor will they ask you to switch products.

Important Safety Information (cont'd)

Talk to your healthcare professional before receiving any vaccinations if you are taking a corticosteroid.

Talk to your doctor about any medications you plan to take including over the counter medications, herbal supplements, and vitamins as certain substances can affect the liver and may reduce the effectiveness of BEQVEZ.

Your healthcare professional will test your factor IX activity levels and for factor IX inhibitors.

After receiving BEQVEZ, your doctor will discuss whether and when you are able to stop prophylaxis, if you need to resume prophylaxis, and actions you may need to take for surgeries, procedures, injuries, and bleeding events.

Do not donate blood, organs, tissues, or cells for transplantation following administration of BEQVEZ.

BEQVEZ is not intended for administration to women. **Males should not donate sperm and should use a male condom or not have sexual intercourse** for up to 6 months after receiving BEQVEZ.

Patients and caregivers should ensure proper handling of any materials that have come into contact with the patient's urine, feces, saliva, mucus, or semen in the first 6 months after BEQVEZ infusion.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see Important Safety Information throughout and full [Prescribing Information](https://www.BEQVEZ.com) at www.BEQVEZ.com.



© 2024 Pfizer Inc. BEQVEZ is a trademark of Pfizer Inc.
All rights reserved. September 2024 PP-BEQ-USA-0427


BEQVEZ[™]
fidanacogene elaparvovec-dzkt
Suspension for intravenous infusion 1 • 10¹¹ vg/mL